

レジメン名【BEP】

1コース:21日間

| [商品名(一般名)] | 投与量 | 1 | 2 | 3 | 4 | 5 | | 8 | | 15 | | 21 |
|------------------|----------------------|---|---|---|---|---|------|---|------|----|------|----|
| ブレオ (ブレオマイシン) | 30mg/body | ● | | | | | | ● | | ● | | |
| ベプシド (エトポシド) | 100mg/m ² | ● | ● | ● | ● | ● | | | | | | |
| シスプラチン | 20mg/m ² | ● | ● | ● | ● | ● | | | | | | |

day1

| | | | | |
|--|--|---|--|--------|
| 8:00-10:00 (250mL/hrで2hr) ソルデム1輸液 500mL | 10:00-12:00 (250mL/hrで2hr) ソルデム1輸液 500mL | 12:00-14:00 (250mL/hrで2hr) ソルアセトF輸液 500mL | 14:00-16:00 (250mL/hrで2hr) ソルアセトF輸液 500mL | 生食シリンジ |
| 9:00-9:30 (100mL/hrで30min) アロキシ0.75mg/バッグ デキサート注射液9.9mg | 10:00-12:00(約150mL/hrで2hr) CDDP 20mg/m ² 生食 250mL | 12:00-14:00(約250mL/hrで2hr) ベプシド 100mg/m ² 生食 500mL | 14:00-14:30 ブレオ 30mg 生食 100mL | |
| 9:15-10:00 (400mL/hrで45min) 20%マンニトール注射液 300mL | | | | |

day2-5

| | | | | |
|---|--|---|--|--------|
| 8:00-10:00 (250mL/hrで2hr) ソルデム1輸液 500mL | 10:00-12:00 (250mL/hrで2hr) ソルデム1輸液 500mL | 12:00-14:00 (250mL/hrで2hr) ソルアセトF輸液 500mL | 14:00-16:00 (250mL/hrで2hr) ソルアセトF輸液 500mL | 生食シリンジ |
| 9:00-9:30 (100mL/hrで30min) デキサート注射液6.6mg 生食 50mL | 10:00-12:00(約150mL/hrで2hr) CDDP 20mg/m ² 生食 250mL | 12:00-14:00(約250mL/hrで2hr) ベプシド 100mg/m ² 生食 500mL | | |
| 9:15-10:00 (400mL/hrで45min) 20%マンニトール注射液 300mL | | | | |

day8 & 15

| |
|----------------------------------|
| (250mL/hrで2hr) ソルアセトF輸液 500mL |
| 30分 ブレオ 30mg 生食 100mL |